

# *In vitro* $\alpha$ -Amylase Inhibitory Activity of the Essential Oils of *Pinus nigra* ARN: A Preliminary Assay

Faten Mezni\*, Sondes Fkiri, M. Larbi Khouja, Zouhair Nasr and Abdelhamid Khaldi

Laboratory of Management and Valorization of Forest Resources, National Researches Institute of Rural engineering, Water and Forests (INRGREF), University of Carthage, Rue Hedi Karray, BP 10 2080-Ariana, Tunisia

## \*Correspondence to:

Faten Mezni

Laboratory of Management and Valorization of Forest Resources, National Researches Institute of Rural engineering, Water and Forests (INRGREF) University of Carthage, Rue Hedi Karray BP 10 2080-Ariana, Tunisia

E-mail: [faten-mez@hotmail.com](mailto:faten-mez@hotmail.com)

Received: February 01, 2022

Accepted: March 09, 2022

Published: March 11, 2022

**Citation:** Mezni F, Fkiri S, Khouja ML, Nasr Z, Khaldi A. 2022. *In vitro*  $\alpha$ -Amylase Inhibitory Activity of the essential oils of *Pinus nigra* ARN.: A Preliminary Assay. *J Food Chem Nanotechnol* 8(1): 18-20.

**Copyright:** © 2022 Mezni et al. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY) (<http://creativecommons.org/licenses/by/4.0/>) which permits commercial use, including reproduction, adaptation, and distribution of the article provided the original author and source are credited.

Published by United Scientific Group

## Abstract

Such as several plants, Pines are known to have anti-diabetic potential. The current study evaluated the anti-diabetic potential of *Pinus nigra* essential oils, obtained from nine *Pinus nigra* provenances planted in Northwestern Tunisia, via *in vitro* inhibition of  $\alpha$ -amylase. The  $\alpha$ -amylase inhibitory potentials were determined using radial diffusion method. Our results showed a significant difference between provenances. The  $\alpha$ -amylase inhibitory potential ranged from 10% to 26.6%. The essential oils of Crimée provenance from *Pinus nigra pallasiana* subsp. displayed the most effective inhibition of  $\alpha$  amylase. While, essential oils of Bois frerot provenance from *Pinus nigra laricio* subsp. showed the lowest inhibitory potential (10%). The observed inhibitions of  $\alpha$ -amylase suggest that the essential oils of *P. nigra* of some provenances may be useful in the management of diabetes mellitus.

## Keywords

*Pinus nigra*, Subspecies, Essential oils, Provenances,  $\alpha$ -Amylase inhibitory activity

## Introduction

Diabetes is a metabolic complex disorder which can be the cause of severe health problems such as hypo or hyper glycemic and coma. It is, sometimes, accompanied by different other complications which can be further difficult to treat than diabetes. Actually, diabetic mellitus is treated by hypoglycemic drugs presenting, almost several side effects. Regarding the large increase in the number of diabetics in the entire world, many researchers have treated the pharmacological effect of several plants [1, 2]. Pines are considered, according to the literature, among the plants that have shown significant anti-diabetic power. Capetti et al. [3] determined the  $\alpha$ -amylase inhibitory potential of essential oils from needles of *Pinus mugo* and *Pinus sylvestris*. In the same context Yang et al. [4] investigated the  $\alpha$ -amylase inhibitory activity of essential oils extracted from the needle of *Pinus tabulaeformis*. To the best of our knowledge, no study was conducted on *Pinus nigra* essential oils. The present investigation evaluated, for the first time, the anti-diabetic potential of essential oils, obtained from nine *Pinus nigra* provenances growing in comparative plantation in Northwestern Tunisia, via *in vitro* inhibition of  $\alpha$ -amylase.

## Materials and Methods

### Plant material

Needles of nine samples of *Pinus nigra* (Table 1) were collected from Souiniet arboretum located in West-Northern Tunisia (8 ° 48 'E, 35 ° 54' N,

**Table 1:** Geographic origin of the nine provenances of *Pinus nigra*.

Sub-species	Provenances	Country	Code
<i>Pinus nigra laricio</i> var. <i>calabrica</i>	Cosenza	Italy	P 1
<i>Pinus nigra laricio</i> var. <i>corsicana</i>	Bois frerot (Ardennes)	France	P 2
<i>Pinus nigra nigra</i> var. <i>austriaca</i>	Puget-Théniers	France	P 3
<i>Pinus nigra nigra</i> var. <i>nigricans</i>	Kustendil	Bulgaria	P 4
<i>Pinus nigra laricio</i> var. <i>calabrica</i>	Cantanzaro	Italy	P 5
<i>Pinus nigra laricio</i> var. <i>corsicana</i>	les Barres (leint)	France	P 6
<i>Pinus nigra pallasiana</i>	Crimée	Russia	P 7
<i>Pinus nigra laricio</i> var. <i>calabrica</i>	Tavola	Italy	P 8
<i>Pinus nigra salzmanni</i>	Olette (Pyr-Orient)	France	P 9

492 m). The plant was identified by Pr. Abdelhamid Khaldi, a specialist in plant identification. The nine samples correspond to provenances from different Mediterranean regions, which have been planted in the North West of Tunisia in 1966.

### Oil extraction

Needles were air-dried and then used to extract essential oil by hydrodistillation. A Clevenger apparatus was used according to the method described by the British Pharmacopoeia [5]. The essential oils were then stored at 4 °C until analysis.

### Oil yield

Oil yield was calculated as the percentage (%) of grams of extracted essential oil per gram of used dry needles.

### $\alpha$ -amylase inhibitory test

The  $\alpha$ -amylase inhibitory potentials were determined using radial diffusion method. 0.1786 g of  $\alpha$ -amylase from *Aspergillus oryzae* was dissolved in 1 mL of 20 mM phosphate buffer and 6 mM of NaCl. The pH value was adjusted to 6.9. To each plate, containing 25 mL of agar-starch, 20  $\mu$ L of  $\alpha$ -amylase solution mixed with 10  $\mu$ L of essential oil were inoculated. Clear diffusion zones were formed after incubation at 20 °C for 4 hours. The percentage of inhibition was calculated according to the following formula:

$$I = \frac{DE - DC}{DC} \times 100$$

Where: DE: Diameter of extract and DC: Diameter of Control

$\alpha$ -amylase inhibitor of wheat seeds (*Triticum aestivum*) was used as the positive control. Results are expressed as the mean of three replications.

### Statistical analysis

The General Linear Models procedure (GLM) of the SAS (9.0) program was used for the statistical analysis.

## Results

Significant variability was determined between the nine provenance ( $p < 0.0001$ ). The most important yield was

recorded by needles of Puget Théniers provenance ( $0.68\% \pm 0.05$ ), however the lowest one was found by les Barres (leint) of *laricio* subspecies ( $0.26\% \pm 0.04$ ) (Table 2).

Results of  $\alpha$ -amylase inhibitory activity showed that the highest effect was recorded for oils from Crimée (Russia) of *pallasiana* subspecies (26.6%) followed by oils from Cosenza (Italy) and from Olette (France) of both *laricio* and *salzmanni* subspecies, respectively (25%) (Figure 1). This inhibition is slightly lower than that of the positive control (35%).

The lowest inhibitory effect was reached by oils from Bois frerot (France) of *laricio* subspecies (10%).


## Discussion

The present study was conducted to determine the potential of essential oils from nine provenances of *P. nigra* subspecies to inhibit  $\alpha$ -amylase enzyme *in vitro*. Significant inhibition effect was demonstrated between the studied oils extracted from needles from different provenances. This variation can be related to the differences in oil composition demonstrated by Fkiri et al. [6].

The potential of essential oils to inhibit key carbohydrate hydrolyzing enzyme has long been studied. Several studies reported the efficiency of these natural products as antidiabetic compounds [7-9]. The anti-diabetic potential of some *Pinus* species were investigated. Capetti et al. [3] studied the  $\alpha$ -amylase inhibitory potential of essential oils from needles of *Pinus mugo* and *Pinus sylvestris*. According to their findings,

**Table 2:** The yield of *Pinus nigra* essential oil.

Provenance	Oil yield (%)
P 1	$0.37^f \pm 0.02$
P 2	$0.41^e \pm 0.04$
P 3	$0.68^a \pm 0.02$
P 4	$0.38^f \pm 0.02$
P 5	$0.51^c \pm 0.05$
P 6	$0.26^i \pm 0.04$
P 7	$0.41^e \pm 0.01$
P 8	$0.66^b \pm 0.02$
P 9	$0.40^e \pm 0.01$



**Figure 1:** Percentage of  $\alpha$ -amylase inhibition by essential oils of *Pinus nigra*.

no activity was observed for the two plants. Yang et al. [4] determined the  $\alpha$ -amylase inhibitory activity of essential oils extracted from the needle of *Pinus tabulaeformis*. An important activity was recorded. It was attributed to terpenoids which are the main types of compounds in the essential oils. The effect of these compounds was highlighted by several studies [10, 11]. Terpenoids have been shown to be the main compounds of *P. nigra* essential oils [12]. This could explain the  $\alpha$ -amylase inhibitory effect demonstrated in our study. To the best of our knowledge, no study addressed the antidiabetic potential of essential oils from *Pinus nigra*.

## Conclusions

This study deals with the preliminary screening of essential oils from needles of *P. nigra* coming from different provenances and subspecies, in order to evaluate their potential hypoglycemic activity.

The observed inhibitions of  $\alpha$ -amylase suggest that the essential oils of *P. nigra* of some provenances may be useful in the management of diabetes mellitus.

More experiments are required to support and deepen these findings. The active molecules have to be identified and isolated.

## Conflict of Interest

Authors declare no conflict of interest.

## References

- Harrower ADB. 1994. Comparison of efficacy, secondary failure rate, and complications of sulfonylureas. *J Diabetes Complications* 8(4): 201-203. [https://doi.org/10.1016/1056-8727\(94\)90044-2](https://doi.org/10.1016/1056-8727(94)90044-2)
- Rhabasa-Lhoret R, Chiasson JL. 2004. -Glucosidase inhibitors. In: Defronzo RA, Ferrannini E, Keen H, Zimmet P (eds) International Textbook of Diabetes Mellitus. John Wiley & Sons Ltd, UK, pp 901-914. <https://doi.org/10.1002/0470862092.d0612>
- Capetti F, Cagliero C, Marengo A, Bicchi C, Rubiolo P, et al. 2020. Bio-guided fractionation driven by in vitro  $\alpha$ -amylase inhibition assays of essential oils bearing specialized metabolites with potential hypoglycemic activity. *Plants* 9(9): 1242-1250. <https://doi.org/10.3390/plants9091242>
- Yang J, Gu D, Wang M, Kou D, Guo H, et al. 2018. In silico-assisted identification of  $\alpha$ -amylase inhibitor from the needle oil of *Pinus tabulaeformis* Carr. *Ind Crops Prod* 111: 360-363. <https://doi.org/10.1016/j.indcrop.2017.10.047>
- British Pharmacopoeia 1988. [<https://journals.sagepub.com/doi/10.1177/106002808902300525>]
- Fkiri S, Mezni F, Rigane G, Ben Salem R, Ghazghazi H, et al. 2021. Chemotaxonomic study of four subspecies of *Pinus nigra* ARN. grown in common garden based on essential oil composition. *J Food Qual* 2021: 5533531. <https://doi.org/10.1155/2021/5533531>
- Loizzo M, Saab A, Atatti G, Menichini F. 2007. Phytochemical composition and in vitro  $\alpha$ -Amylase inhibitory effect of essential oils from *Cedrus libani*. *Fitoterapia*. 78(4): 323-326. <https://doi.org/10.1016/j.fitote.2007.03.006>
- Tahir HU, Sarfraz RA, Ashraf A, Adil S. 2016. Chemical composition and anti-diabetic activity of essential oils obtained from two spices (*Syzygium aromaticum* and *Cuminum cyminum*). *Inter J Food Prop* 19(10): 2156-2164. <https://doi.org/10.1080/10942912.2015.1110166>
- Ojah EO, Moronkola DO, Akintunde AAM. 2020.  $\alpha$ -amylase and  $\alpha$ -glucosidase antidiabetic potential of ten essential oils from *Calophyllum inophyllum* Linn. *Iberoam J Med* 2(4): 253-260. <http://dx.doi.org/10.5281/zenodo.3841108>
- Jelenkovic L, Jovanovic VS, Palic I, Mitic V, et al. 2014. In vitro Screening of  $\alpha$ -amylase inhibition by selected terpenes from essential oils. *Trop J Pharm Res* 13(9): 1421-1428. <https://doi.org/10.4314/tjpr.v13i9.7>
- Uddin N, Hasan MR, Hossain MM, Sarker A, Hasan AHMN, et al. 2014. In vitro  $\alpha$ -amylase inhibitory activity and in vivo hypoglycemic effect of methanol extract of *Citrus macroptera* Montr. fruit. *Asian Pac J Trop Biomed* 4(6): 473-479. <https://doi.org/10.12980/AP-JTB.4.2014C1173>
- Amri I, Hanana M, Jamoussi B, Hamrouni L. 2017. Essential oils of *Pinus nigra* Arnold subsp. *laricio* Maire: Chemical composition and study of their herbicidal potential. *Arab J Chem* 10(2): 3877-3882. <https://doi.org/10.1016/j.arabjc.2014.05.026>