

Mycotoxins' Contamination in Food and Feeds: A Review

Afsaneh Morshedi¹, Ahmad Shakerardekani², Reza Karazhian³, Mohammad Morshedi⁴, Marcos Eduardo Valdes¹, Toktam Mohammadi-Moghaddam^{5*} and Mohaddeseh Kariminejad⁵

¹Facultad de Ciencias Gastronómicas y Turismo, Universidad UTE, Quito, Ecuador

²Pistachio Research Center, Horticultural Sciences Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Rafsanjan, Iran

³Academic Center for Education, Culture and Research of Iran, Iran

⁴Department of Geophysics, Tehran University, Tehran, Iran

⁵Department of Food Science and Technology, Neyshabur University of Medical Sciences, Neyshabur, Iran

*Correspondence to:

Toktam Mohammadi-Moghaddam
Department of Food Science and Technology,
Neyshabur University of Medical Sciences,
Neyshabur, Iran.

E-mail: mohammadit1@nums.ac.ir

Received: October 03, 2023

Accepted: November 03, 2023

Published: November 06, 2023

Citation: Morshedi A, Shakerardekani A, Karazhian R, Morshedi M, Valdes ME, et al. 2023. Mycotoxins' Contamination in Food and Feeds: A Review. *J Food Chem Nanotechnol* 9(3): 141-149.

Copyright: © 2023 Morshedi et al. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY) (<http://creativecommons.org/licenses/by/4.0/>) which permits commercial use, including reproduction, adaptation, and distribution of the article provided the original author and source are credited.

Published by United Scientific Group

Abstract

Fungi are the most resistant microorganisms distributed worldwide in different foods and feedstuffs. They produce Mycotoxins as their secondary and toxic metabolites for humans and animals, and they are carcinogenic, mutagenic, teratogenic, and immunosuppressive. According to Mycotoxins structures, their risk for health is different. Therefore, aflatoxins, ochratoxins A, zearalenone, fumonisins, patulinand, and trichothecenes are recognized as the most hazardous for health. Climate changes during recent years have caused increasing mycotoxins contaminations. Since these secondary metabolites cause chronic and acute diseases, different standards exist to eliminate them in foodstuffs worldwide. The toxicity of mycotoxins and their maximum permitted in foods is different. Because of their toxicity feature, many scientists are studying to find better options for mycotoxin detection and reduction in foods and feeds. Up to now, HPLC (High performance liquid chromatography) coupled with fluorescence derivatization and MS/MS (Mass spectroscopy) is the most successful method for mycotoxin detection. There are different ways to reduce these contaminants, but in countries that have considerable rainfall besides temperature, which helps to grow molds and produce mycotoxins significantly.

Keywords

Aflatoxins, Ochratoxin, Patulin, Fumonisins, Zearalenone

Introduction

Nowadays, we face climate change all over the world. These changes have produced adverse effects on all aspects of food and nutrition security and safety, for instance: high temperature, CO₂ (carbon dioxide) increasing, rainfall amount and its distribution, disease and pest distribution which have led to increasing mycotoxins contaminations [1-8]. In the next 10 - 20 years, the environment will change markedly with atmospheric CO₂ concentration with an increasing rate of 1.5 μmol/year [9]. Not only increasing CO₂ but also greenhouse gases, causes global warming and are expected to increase the rate of air temperature by 0.03 °C per year. Paterson and Lima [10] suggest that if the temperature increases during future years, it will observe those aflatoxins and other mycotoxins production in crops will be increased consequently.

Another critical point that is rarely noticed is that most molds can fetch up several types of mycotoxins at the same time, which can show interaction too. Since food and feed can be substrates of several fungi simultaneously, humans

and animals face high contents of different types of mycotoxins [1, 11].

Among the emerging issue in food safety, is the increase in the occurrence of toxigenic fungi (*Aspergillus*, *Fusarium*, and *Penicillium*) and their metabolism (aflatoxins, ochratoxins, patulin, zearalenone, fumonisin, and deoxynivalenol) is the primary concern because they are carcinogenic, mutagenic, teratogenic, and immunosuppressive. Finally, they are responsible for acute or chronic toxic effects with chronic disease in the central system of nervous, liver, and cardiovascular [12-19]. These contaminations always occur in poor socio-economic situations not only in pre-harvest but also in post-harvest stages, and consequently related health problems will occur with these products. The best point relating to these mycotoxins is that they can affect animals and their products by feeding too, and finally their adverse effects will be seen in humans who consume them [20, 21]. Mycotoxins are very stable, and this feature causes them to make pollution in different agro products, especially cereals, at different stages like pre-harvest, harvest, and post-harvest. Therefore, they are categorized as field and storage fungi [22-24]. Since these toxins production depends on climate and environmental conditions, product contaminations will vary in different countries [25, 26]. Different environmental factors, for instance, temperature, mold species, substrate, moisture percent, relative humidity, water activity, airflow, and ambient brightness, can affect toxin production. Besides some physical and chemical parameters like insect damages, fungicides application, storage conditions, poor operation during transportation and storage, and inadequate ventilation mycotoxins production can be different [25, 27].

Up to 1985 FAO (Food and Agriculture Organization) assessed that 25% of global food crops contaminated by mycotoxins, but today the EU (European Union) estimated to be up to 60 - 80%. Of course, this high occurrence results from improved analytical methods and climate change during these years, by the way it is a global problem [28]. Approximately 25 - 50% of cereal and its products are mycotoxins contaminated worldwide, and 5 - 10% is not consumable and produces considerable economic loss [7, 29, 30].

There are many identified mycotoxins but the most population of them are aflatoxins, ochratoxins A, zearalenone, fumonisin, patulin, and trichothecenes [31]. Mycotoxins can be produced in the field, garden, or farm, and by polluting the raw materials can be conducted to processed food [24, 32]. While the first death reason in the world has been reported is a chronic disease, like cardiovascular, the second reason is respiratory diseases, and the third is all types of cancers; mycotoxin poisonings stand after them [33-37]. In table 1, different mycotoxins effects on humans are mentioned. Therefore, these secondary metabolites are responsible for major public health that governments should be aware of as much as air pollution. The FAO estimated that more than 25% of agro-foods worldwide are contaminated by mycotoxins and 4.5 - 5 billion people are at risk of chronic exposure to mycotoxins especially in poor countries [38, 39].

Different studies have demonstrated that the most mycotoxin contaminations belong to the parts with hot and hu-

mid climates, tropical, and Estevan regions. Mycotoxins are separated from a wide range of agro-products: cereals (wheat, rice, maize, barley, etc.); nuts (pistachio, peanut, almond, etc.); spices (pepper, cayenne, cinnamon, etc.); fruits and vegetables (apple, cherry, etc.); animal products (meat, milk, etc.) [40-46]. In figure 1, different factors for mycotoxin occurrence in food and feed are shown.

Mycotoxins consumption and intake lead to chronic illnesses: immune sickness [47], metabolic, biochemical, and allergic illnesses [48, 49], carcinogenicity, teratogenicity, mutagenicity, and death. Therefore, because of the high possibility of their intake of food and agricultural products, it is necessary for their content management in foods and even feeds [50-52]. Mycotoxins are entirely different in physicochemical properties and follow different pathways after intake. Since their body effect and toxicity is completely different, they categorized according to IARC (International Agency for Research on Cancer) to different groups (Table 2).

Types of Mycotoxins

Aflatoxins

These toxins are products of *Aspergillus* species through a polypeptide pathway [53]. They are mostly observed in nuts and cereals. In most cereal crops, these contaminations occur after harvest and during storage. Scientists believe high temperatures and relative humidity more than 65% are the profit situation for these toxins production [54].

Aflatoxin's structure

Six types of naturally existing aflatoxins are aflatoxin B₁, aflatoxin B₂, aflatoxin G₁, aflatoxin G₂, aflatoxin M₁, and aflatoxin M₂. Aflatoxins M₁ and M₂ are recognized in milk for the first time; therefore, the letter (M) comes from milk. Their origin is AFB₁ after animal body intake, restructure occurs and

Table 1: Mycotoxins effects on different humans' organs [39].

Mycotoxins	Humans' organs
Aflatoxins	Liver, small intestine, lymph nodes
Fumonisin	Brain, lymph nodes, stomach
Ochratoxin A	Kidney
Patulin	Lymph nodes
Sterigmatocystin	Lymph nodes, windpipe and throat




Figure 1: Different factors for mycotoxin occurrence in food and feed [29, 45, 46].

Table 2: Foods and different mycotoxins contaminations.

Group	Mycotoxin	Ref.
Cereals	Trichothecenes	[86]
	Aflatoxins	JECFA 1997, 1999, 2012, 2017; EFSA 2007, 2004a; IARC 2015, WHO 2000
	Fumonisin	IARC 2015; JECFA 2012, 2017; EFSA 2018a
	Deoxynivalenol	EFSA 2004b, 2011, 2017d; JECFA 2001, 2011
	Zearalenones	EFSA 2011, 2014, 2016, 2017c; JECFA 2000
	Ochratoxin	ESFA 2006; IARC 1983, 1993; JECFA 1996, 2001, 2007, 2008
Spices	Ochratoxin	ESFA 2006; IARC 1983, 1993; JECFA 1996, 2001, 2007, 2008
Nuts	Aflatoxins	JECFA 1997, 1999, 2012, 2015, 2017; IARC 2002, 2015; EFSA 2004a, 2007; WHO 2000
	Ochratoxins	ESFA 2006; IARC 1983, 1993; JECFA 1996, 2001, 2007, 2008
Fruits and vegetables	Patulin	-
	Ochratoxins	EFSA 2006; IARC 1983, 1993; JECFA 1996, 2001, 2007, 2008
Coffee and cacao	Ochratoxins	EFSA 2006; IARC 1983, 1993; JECFA 1996, 2001, 2007, 2008
Milk	Aflatoxins	JECFA 1997, 1999, 2017; IARC 2002; EFSA 2004a, 2007
	Fumonisin	IARC 2015; JECFA 2012, 2017
	Zearalenones	EFSA 2011, 2014, 2016, 2017c; IARC 1993; JECFA 2000; SCF 200a
	Ochratoxins	EFSA 2006; IARC 1983, 1993; JECFA 1996, 2001, 2007, 2008
Egg	Zearalenones	-
	Ochratoxins	EFSA 2006; IARC 1983, 1993; JECFA 1996, 2001, 2007, 2008

aflatoxin M₁ and aflatoxin M₂ will be formed [28]. The other four of these different types of aflatoxins are named because of their colors under ultraviolet shining during thin layer chromatography (TLC) analysis; B (Blue) and G (Green). On the thin layer, these mentioned aflatoxins have different retention factors. That's means toxins with the number (1) index have more relative mobility on TLC, number (2) index shows less mobility [28, 55].

Aflatoxins toxicity

Aflatoxins are the most important mycotoxins because of their genotoxic carcinogenic features; they are the most potent mutagenic and carcinogenic materials [28]. Toxicity rates are different in these types of aflatoxins: B₁ > G₁ > M₁ > B₂ > G₂ > M₂ [55]. The IARC recognized that AFB₁ is the most hazardous mycotoxin for human health [28, 56]. The severity of disease after aflatoxins consumption depends on the amount and duration of eating the toxin. Vomiting, hemorrhage, abdominal pain, icterus, cerebral edema, coma, convulsions, pulmonary and even death are the symptoms of aflatoxicosis which is aflatoxins poisoning disease. Besides the symptoms it is necessary to be mentioned that aflatoxicosis in chronic position can be observed as tumors and cancers especially liver cancer [28]. Because of these health problems produced by aflatoxins, United States Food and Drug Administration (USFDA) and EU, have set the maximum level for aflatoxins contamination in foods which are less than 20 µg/kg, and 0.1 - 2 µg/kg respectively for total aflatoxins [57]. The important point is that Sterigmatocystin is a genotoxic carcinogen. Its biosynthesis pathway is the same as aflatoxins, and finally both have the same target organ (liver) [58].

Fumonisin

These types of toxins are second metabolites of *Fusarium* species in fields, which produced in hot and dry or humid weather [59]. Up to now, more than 30 different types of fumonisins have been recognized while their number is growing [60]. Fumonisin were seen for the first time in cereal, especially maize (corn) [61, 62].

Fumonisin structure

Fumonisin have polyketide non-fluorescent structures. Fumonisin considering their chemical structures is divided into four categories: A, B, C, and P [63]. In fumonisin B group (fumonisin B₁, fumonisin B₂, fumonisin B₃, and fumonisin B₄), fumonisin is the most prevalent and toxic one and is observed 70 - 80% more than the others. Fumonisin B₂ and fumonisin B₃ co-occur with fumonisin B₁, but among the fumonisin Bs group, fumonisin B₁ is considered in Group 2B possibly carcinogenic to humans by the IARC [56, 58].

Fumonisin toxicity

The reported symptoms are disruption of sphingolipids metabolism and effecting on neural tube defects in the brain and spinal, which are very similar to sphingosine and produce some neurological manifestations such as nervousness, ataxia, aimless circling, lameness, facial paralysis, abnormal movements and finally inability to drink or eat. However, the target organ for fumonisin B in laboratory animals is the liver. Fumonisin B₁ is categorized in IARC 2B which means it is possibly carcinogenic to humans [64]. According to FDA standards, the maximum advisory level of fumonisin Bs is 2 - 4 mg/kg for human consumption.

Ochratoxins

Ochratoxins are another mycotoxins group that can be produced by *Aspergillus ochraceous*, *Aspergillus carbonarius*, and *Penicillium verrucosum* species mostly [65]. It is separated from different foods (e.g., cereals, coffee and cacao beans, nuts, spices, milk, and egg) [66].

Ochratoxin structure

There are different functional groups in this toxin main structure therefore, according to that, ochratoxins can be exported into three groups: A, B, and C. The phenylalanyl derivative of isocoumarin substituted is named ochratoxin A which is recognized as the second most important mycotoxin [67]. The pathway of ochratoxins A is not known completely against aflatoxins and fumonisin Bs, but it recognized isocoumarin group is originated from acetate and molanate via a polyketide synthesis pathway [68].

Ochratoxin toxicity

When ochratoxins A enters the body through foods, it binds to plasma protein through gastrointestinal and sediment in the kidney with a very long half-life (about 35 days) [28]. Besides, ochratoxins A can corral with phenylalanine hydroxide in both kidney and liver to stop certain protein synthesis as well as ribonucleic acid (RNA) and deoxyribonucleic acid (DNA) synthesis. Ochratoxins A has been recognized as a carcinogen factor in animal studies therefore, it categorized in 2B group possibly carcinogenic to humans by the IARC [56]. The FDA has not set regulations on ochratoxins A content in foods, while the EU has established ochratoxins A limitations of 3 and 5 mg/kg for food products and raw, respectively.

Deoxynivalenol

Deoxynivalenols are mainly produced by *Fusarium graminearum* and *Fusarium culmorum* in the field or during post-harvest storage mainly in cereals [28].

Deoxynivalenol structure

More than 200 different trichothecenes have been recognized up to know that is subdivided into four main groups: A, B, C, and D. Type A of them, is known as the most toxic group as for example T-2 and HT-2 are in this category [5]. Chemically, deoxynivalenol is considered a type B of trichothecene with a characteristic ketone at C8, and three hydroxyl groups at C3, C7, and C15. Another derives of deoxynivalenol are its acetylated, 3-acetyldeoxynivalenol (3ADON), and nivalenol (NIV) which is type B trichothecenes and deoxynivalenol-3-glucoside [25, 69]. According to some research, deoxynivalenols have an amphipathic nature; therefore, they can cross the cell membrane and interact with endoplasmic reticulum, mitochondria [25], and chloroplast [25, 70].

Deoxynivalenol toxicity

The epoxide ring of trichothecenes has toxicity action that its solution is not only stable against PH but also against heat [71]. According to [72, 73] studies, deoxynivalenols effects can inhibit cell dividing, synthesis of RNA and DNA, membrane disruption, and mitochondrial functions. Deoxynivalenols prevent initial peptide bonds. Meanwhile, although

deoxynivalenols inhibit DNA and protein synthesis, there is no report about its mutagenic or carcinogenic effect in humans. They affect immunity by protein synthesis inhibition [67]; decrease antibody and immunoglobulin levels in humans [74]; easily absorbed through skin and cause rapid irritation. In animals, weight loss and anorexia are the most common chronic effects. It confirmed that deoxynivalenol toxicity is less than type 'A' trichothecene. As well, these types of mycotoxins consumption have produced fatal and chronic toxicoses not only in humans but also in animals. They can affect plant regeneration and mammals' reproduction [75]. Deoxynivalenol poisonous symptoms are vomiting, diarrhea, headache, dizziness, abdominal pain, and fever [76]. These family mycotoxins are recognized as phytotoxic for parsnip, wheat, and maize. The flowering period is considered a critical stage for plants for *Fusarium* species. Contaminants that can make progress in warm and moist conditions. T-2 and HT-2 chronic consumption symptoms are hematotoxicity and immunotoxicity and in acute situations vomiting in animals [77, 78]. The IARC categorized deoxynivalenol and T-2 toxin in group 3 which are not carcinogenic to humans. According to FDA standards, the maximum content of deoxynivalenol should be 1 mg/kg in final food products [56].

Zearalenone

Zearalenone is a common contamination in maize and cereal grains and products. A field fungus that grows in cool and moist conditions during different stages: pre-harvest, post-harvest, and storage. However, toxin production usually happens after harvesting and during storage and it is observed in cereals [79]. The zearalenone name was F-2 toxin previously [80] and is achieved from polyketide pathway by *Fusarium* species [81]. Another interesting point is that cereal contamination by zearalenone and deoxynivalenol usually coincide simultaneously because the same fungus produces both [82].

Zearalenone structure

The structure of zearalenone is (3, 4, 5, 6, 9, 10-hexahydro-14, 16-dihydroxy-3-methyl-1H-2 benzoxacyclotetradecin-1, 7(8H)-dione. Therefore, it is a macro cyclic β -resorcylic acid lactone [67]. Because of zearalenone complicated chemical structure, it is a heat resistant mycotoxin. To degrade it, should use alkaline solutions combine a temperature higher than 150 °C. However, children who consume cereal-based foods might receive this toxin more than others [25, 67].

Zearalenone toxicity

The IARC data shows that zearalenone is classified as group 3 and is not a carcinogen for humans [56]. The maximum zearalenone level in human foods depends on the food category and should be 75 - 350 μ g/kg, according to EU standards. The FDA defines no limitation [68, 83].

Patulin

Patulin is the second metabolite of 60 different species of *Penicillium* which is considered a serious hazard for fruits, especially during the post-harvest period [5, 29]. The mold first grows on the surface of the fruit, like apples, cherries, figs, etc., and then contaminates all the fruit and fruit juice [29, 84, 85].

The reported data has shown that approximately 50% of apple juice is highly contaminated by patulin worldwide [86].

Patulin structure

Structurally patulin is a heterocyclic lactone (4-hydroxi-4H-furo(3, 2-c)piran-2(6H)-ona) [5]. It is a water-soluble and colorless molecule.

Patulin toxicity

It is supposed to cause patulin toxicity in its reaction with thiol groups (cysteine, glutathione, etc.) in the cytoplasm [87]. It affects enzymes that usually are digestion, metabolism, and energy production responsible [87]. Therefore, patulin chronic toxicity symptoms are nausea, vomiting, gastrointestinal, kidney and liver damage, and immunosuppression in animals [29]. It is not clear that patulin is a carcinogen for humans [5]. Both EU and FDA have set an upper limit of 50 µg/L or µg/kg patulin in apple and fruit juices, while for solid apple products like compote or puree, it is 25 µg/kg [5, 40, 88].

Mycotoxins Analysis

All the analysis methods have different risks that can be classified into five categories: health, safety, environment, energy, and waste based on toxicity [89].

For mycotoxin analysis, the method should be quick, cheap, and effective besides accurate detection. Aflatoxins, ochratoxins, fumonisins, and zearalenones have similar chemical structures with the same physicochemical properties [68, 90-92]. Due to the use of hazardous and toxic chemical materials, the analysis should be done according to set burdens. Nowadays, green analysis is developed by different research societies: Analytical Eco-Scales [93], National Environmental Methods Index, and Green Analytical Procedure Index [89]. All these methods are planned to reduce environmental damage and take into rapid and efficient analysis [94]. According to different researcher's publications, there are routine and different stages for mycotoxin analysis:

Sample collection

The first step of every analytical procedure is sample gathering. There are four ways for this step: in-line, on-line, at-line, and off-line sampling [89] of these models, the greenest one is in-line sampling, online and at-line are categorized in medium-green approach and off-line sampling should be avoided [95].

How many should be a sample size? It completely depends on the bulk weight and food type [96]. For example, if we have 50,000 kg of cereals, we should gather at least 100 incremental samples and the final weight should be 10 kg. There is a rule for sampling according to 'lot' and 'sublot' weight.

The next important step related to sample gathering is sample protection until analysis. There are three methods of preservation: chemical, physical, and physicochemical. Choosing each method is wholly dependent on the sample nature [89].

Extraction procedure

This procedure is the most crucial stage in every analyti-

cal experiment. In this step, considering the physicochemical properties of the sample (food matrix) and the mycotoxin that should be analyzed, the extraction method is different. However, grinding, homogenization with organic solvents or strong acids, and finally, filtration are its steps [89, 96].

The routine methods of these steps are liquid-liquid solvent extraction, immuno-affinity column extraction, solid-phase extraction, and the newest one is pressurized hot water extraction [97]. During extraction, the solvent will be removed analyte from the food matrix. Certainly, solvent choosing has a key role in this procedure and the point is that there is no specific solvent for an analyte or food. The best solvent is the solvent that removes as much analyte as possible from the food matrix [96].

Most of the mycotoxins are soluble in polar solvents. Therefore, solvents, used in the extraction process, up to now, are water, acetone, acetonitrile, methanol, chloroform, potassium chloride, or a combination of these [96]. Solvent and ground food should mix completely by mechanical shaking, ultrasound, supercritical fluid extraction, accelerated solvent extraction, and microwave-assisted extraction. Finally, the eluate will evaporate under the nitrogen stream. The following steps are re-dissolution and filtration [90-92, 94, 96].

Purification (clean up) procedure

The aim of purification is to separate target molecules or molecules. Each sample extract contains co-extracted materials that interfere with the analyte during detection with analytical instruments.

Solid phase extraction (SPE) is based on molecularly imprinted polymers and used for this purpose. These columns are different and produced commercially. According to matrix, analyte, and interference, it is necessary to choose the column. The newest and best purification method is using immune-affinity column (IAC). The point is that these columns act completely specific and for each type of mycotoxin, it is needed to use specified columns because they contain antibodies [94, 96]. In some of the analyses, it recommended to use both SPE and IAC for purification [96].

Detection and quantification

Analytical instrument selection is the main factor in the success of an analytical method. There are official mycotoxins methods approved by regulatory authorities such as the USFDA, The Association of Official Analytical Chemists, and the European Commission. The base of all official methods is the chromatography method. Commonly, Liquid Chromatography (HPLC or UHPLC) coupled to the tandem mass spectrometer (MS/MS) is recognized as the industry standard for mycotoxins analysis.

Among chromatographic techniques, HPLC coupled with fluorescence derivatives and MS/MS is the most popular method for cereal mycotoxin detection [98]. Some scientists are studying new methods focusing on multi-mycotoxin extraction, which not only have lower costs but are also categorized as green methods (like LC-MS/MS), but also is not an official method yet [97].

Conclusion

Mycotoxins are harmful second metabolites of fungi with different toxicity. The most harmful toxicity observed in aflatoxins fungi contaminants start from the field and continue during transportation, storage, and manufacturing; finally, they will transport to food and feed. During this way, many direct factors like crops genetics, instability of toxigenic properties, fungal species are important. Other factors like climate changes (CO₂ and O₂ concentration, temperature, and relative humidity), moisture, mechanical injuries, pesticides, and fungicides are important as well. Mycotoxins can be transported to humans directly and indirectly. When a person eats mycotoxin contaminated food, they directly consume it, but indirectly he/she can intake it by animal products e.g., milk, egg, meat, etc. Therefore, the best way for prevention consuming mycotoxins is molds growing control in different part of food and feed production chain.

Acknowledgements

None.

Conflict of Interest

The authors declare that they do not have any conflict of interest.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- Grenier B, Oswald I. 2011. Mycotoxin co-contamination of food and feed: meta-analysis of publications describing toxicological interactions. *World Mycotoxin J* 4(3): 285-313. <https://doi.org/10.3920/WMJ2011.1281>
- World Agriculture: Towards 2030/2050 - The 2012 Revision. [<https://www.fao.org/global-perspectives-studies/resources/detail/en/c/411108/>] [Accessed November 05, 2023]
- Medina A, Akbar A, Baazeem A, Rodriguez A, Magan N. 2017. Climate change, food security and mycotoxins: do we know enough?. *Fungal Biol Rev* 31(3): 143-154. <https://doi.org/10.1016/j.fbr.2017.04.002>
- Firdaus RR, Gunaratne MS, Rahmat SR, Kamsi NS. 2019. Does climate change only affect food availability? What else matters? *Cogent Food Agric* 5(1): 1707607. <https://doi.org/10.1080/23311932.2019.1707607>
- Agriopoulou S, Stamatelopoulou E, Varzakas T. 2020. Advances in occurrence, importance, and mycotoxin control strategies: prevention and detoxification in foods. *Foods* 9(2): 137. <https://doi.org/10.3390/foods9020137>
- Conte G, Fontanelli M, Galli F, Cotrozzi L, Pagni L, et al. 2020. Mycotoxins in feed and food and the role of ozone in their detoxification and degradation: an update. *Toxins* 12(8): 486. <https://doi.org/10.3390/toxins12080486>
- Aasa AO, Fru FF, Adelusi OA, Oyeyinka SA, Njobeh PB. 2023. A review of toxigenic fungi and mycotoxins in feeds and food commodities in West Africa. *World Mycotoxin J* 16(1): 33-47. <https://doi.org/10.3920/WMJ2021.2766>
- Truong NN, Tesfamariam K, Visintin L, Goessens T, De Saeger S, et al. 2023. Associating multiple mycotoxin exposure and health outcomes: current statistical approaches and challenges. *World Mycotoxin J* 16(1): 25-32. <https://doi.org/10.3920/WMJ2022.2784>
- Magan N, Medina A, Aldred D. 2011. Possible climate-change effects on mycotoxin contamination of food crops pre-and postharvest. *Plant Pathol* 60(1): 150-163. <https://doi.org/10.1111/j.1365-3059.2010.02412.x>
- Paterson RRM, Lima N. 2010. How will climate change affect mycotoxins in food? *Food Res Int* 43(7): 1902-1914. <https://doi.org/10.1016/j.foodres.2009.07.010>
- Xu F, Baker RC, Whitaker TB, Luo H, Zhao Y, et al. 2022. Review of good agricultural practices for smallholder maize farmers to minimise aflatoxin contamination. *World Mycotoxin J* 15(2): 171-186. <https://doi.org/10.3920/WMJ2021.2685>
- Kebede H, Liu X, Jin J, Xing F. 2020. Current status of major mycotoxins contamination in food and feed in Africa. *Food Control* 110: 106975. <https://doi.org/10.1016/j.foodcont.2019.106975>
- Stroka J, Maragos CM. 2016. Challenges in the analysis of multiple mycotoxins. *World Mycotoxin J* 9(5): 847-861. <https://doi.org/10.3920/WMJ2016.2038>
- Han X, Huangfu B, Xu T, Xu W, Asakiya C, et al. 2022. Research progress of safety of zearalenone: a review. *Toxins* 14(6): 386. <https://doi.org/10.3390/toxins14060386>
- Yu H, Zhang J, Chen Y, Zhu J. 2022. Zearalenone and its masked forms in cereals and cereal-derived products: a review of the characteristics, incidence, and fate in food processing. *J Fungi* 8(9): 976. <https://doi.org/10.3390/jof8090976>
- Ekwomadu T, Mwanza M, Musekiwa A. 2022. Mycotoxin-linked mutations and cancer risk: a global health issue. *Int J Environ Res Public Health* 19(13): 7754. <https://doi.org/10.3390/ijerph19137754>
- Ghazi T, Arumugam T, Foolchand A, Chaturgoon AA. 2020. The impact of natural dietary compounds and food-borne mycotoxins on DNA methylation and cancer. *Cells* 9(9): 2004. <https://doi.org/10.3390/cells9092004>
- Malvandi AM, Shahba S, Mehrzad J, Lombardi G. 2022. Metabolic disruption by naturally occurring mycotoxins in circulation: a focus on vascular and bone homeostasis dysfunction. *Front Nutr* 9: 915681. <https://doi.org/10.3389/fnut.2022.915681>
- Wang K, Liu S, Zhang X, Sun D. 2023. Toxic effect of mycotoxins on cardiovascular system: a topic worthy of further study. *Food Rev Int* 39(4): 2203-2211. <https://doi.org/10.1080/87559129.2021.1950172>
- Wild CP, Gong YY. 2010. Mycotoxins and human disease: a largely ignored global health issue. *Carcinogenesis* 31(1): 71-82. <https://doi.org/10.1093/carcin/bgp264>
- Ozturkoglu-Budak S, Akal HC, Öztürk Hİ. 2023. Multi-mycotoxin production of cheese-derived fungal strains *in vitro* and in cheese models. *World Mycotoxin J* 1-12. <https://doi.org/10.3920/WMJ2023.2831>
- Nleya N, Adetunji MC, Mwanza M. 2018. Current status of mycotoxin contamination of food commodities in Zimbabwe. *Toxins* 10(5): 89. <https://doi.org/10.3390/toxins10050089>
- Mohammed A, Seyoum C, Yousuf J, Mweetwa A, Odera JA, et al. 2023. Multi-mycotoxins analysis in post-harvest maize (*Zea mays* L.) grain from major producing areas of Ethiopia. *World Mycotoxin J* 1-12. <https://doi.org/10.3920/WMJ2022.2829>
- Getahun M, Fininsa C, Mohammed A, Bekeko Z, Sulyok M. 2023. Fungal species and multi-mycotoxin in bread wheat (*Triticum aestivum* L.) in Ethiopia. *World Mycotoxin J* 16(2): 179-194. <https://doi.org/10.3920/WMJ2022.2820>
- Afsah-Hejri L, Jinap S, Hajeb P, Radu S, Shakibzadeh SH. 2013. A review on mycotoxins in food and feed: Malaysia case study. *Compr Rev Food Sci Food Saf* 12(6): 629-651. <https://doi.org/10.1111/1541-4337.12029>
- Mahuku G, Mauro A, Pallangyo B, Nsami E, Boni SB, et al. 2023. Atoxicogenic-based technology for biocontrol of aflatoxin in maize and

- groundnuts for Tanzania. *World Mycotoxin J* 16(1): 59-73. <https://doi.org/10.3920/WMJ2021.2758>
27. Darwish WS, Ikenaka Y, Nakayama SM, Ishizuka M. 2014. An overview on mycotoxin contamination of foods in Africa. *J Vet Med Sci* 76(6): 789-797. <https://doi.org/10.1292/jvms.13-0563>
 28. Eskola M, Kos G, Elliott CT, Hajšlová J, Mayar S, et al. 2020. Worldwide contamination of food-crops with mycotoxins: validity of the widely cited 'FAO estimate' of 25%. *Crit Rev Food Sci Nutr* 60(16): 2773-2789. <https://doi.org/10.1080/10408398.2019.1658570>
 29. Haque MA, Wang Y, Shen Z, Li X, Saleemi MK, et al. 2020. Mycotoxin contamination and control strategy in human, domestic animal and poultry: a review. *Microb Pathog* 142: 104095. <https://doi.org/10.1016/j.micpath.2020.104095>
 30. Wang Y, Zhao C, Zhang D, Zhao M, Zheng D, et al. 2018. Simultaneous degradation of aflatoxin B1 and zearalenone by a microbial consortium. *Toxicon* 146: 69-76. <https://doi.org/10.1016/j.toxicon.2018.04.007>
 31. Udomkun P, Wiredu AN, Nagle M, Bandyopadhyay R, Müller J, et al. 2017. Mycotoxins in Sub-Saharan Africa: present situation, socio-economic impact, awareness, and outlook. *Food Control* 72: 110-122. <https://doi.org/10.1016/j.foodcont.2016.07.039>
 32. Morshedi A, Razavi SMA. 2020. Effect of infrared roasting process on the microorganism contaminations of long and round Iranian pistachio kernels. *J Nuts* 11(1): 23-36. <https://doi.org/10.22034/JON.2020.1869504.1057>
 33. Eze UA, Routledge MN, Okonofua FE, Huntriss J, Gong YY. 2018. Mycotoxin exposure and adverse reproductive health outcomes in Africa: a review. *World Mycotoxin J* 11(3): 321-339. <https://doi.org/10.3920/WMJ2017.2261>
 34. Gibb H, Devleeschauwer B, Bolger PM, Wu F, Ezendam J, et al. 2015. World Health Organization estimates of the global and regional disease burden of four foodborne chemical toxins, 2010: a data synthesis. *PLoS One* 10: e0147996. <https://doi.org/10.1371/journal.pone.0147996>
 35. Wu F. 2015. Global impacts of aflatoxin in maize: trade and human health. *World Mycotoxin J* 8(2): 137-142. <https://doi.org/10.3920/WMJ2014.1737>
 36. Andrade PD, Caldas ED. 2015. Aflatoxins in cereals: worldwide occurrence and dietary risk assessment. *World Mycotoxin J* 8(4): 415-431. <https://doi.org/10.3920/WMJ2014.1847>
 37. Groopman JD, Smith JW, Rivera-Andrade A, Alvarez CS, Kroker-Lobos MF, et al. 2021. Aflatoxin and the aetiology of liver cancer and its implications for Guatemala. *World Mycotoxin J* 14(3): 305-317. <https://doi.org/10.3920/WMJ2020.2641>
 38. Imes T. 2011. The implications of aflatoxin contamination for local food safety in Senegal. USDA Technical Policy Paper.
 39. Braicu C, Buse M, Busuioc C, Drula R, Gulei D, et al. 2019. A comprehensive review on MAPK: a promising therapeutic target in cancer. *Cancers* 11(10): 1618. <https://doi.org/10.3390/cancers11101618>
 40. Vidal A, Mengelers M, Yang S, De Saeger S, De Boevre M. 2018. Mycotoxin biomarkers of exposure: a comprehensive review. *Compr Rev Food Sci Food Saf* 17(5): 1127-1155. <https://doi.org/10.1111/1541-4337.12367>
 41. Wu F, Bhatnagar D, Bui-Klimke T, Carbone I, Hellmich R, et al. 2011. Climate change impacts on mycotoxin risks in US maize. *World Mycotoxin J* 4(1): 79-93. <https://doi.org/10.3920/WMJ2010.1246>
 42. Torres O, Matute J, Gelineau-van Waes J, Maddox JR, Gregory SG, et al. 2015. Human health implications from co-exposure to aflatoxins and fumonisins in maize-based foods in Latin America: Guatemala as a case study. *World Mycotoxin J* 8(2): 143-159. <https://doi.org/10.3920/WMJ2014.1736>
 43. Dong Y, Fan L, Liang J, Wang L, Yuan X, et al. 2020. Risk assessment of mycotoxins in stored maize: case study of Shandong, China. *World Mycotoxin J* 13(2): 313-320. <https://doi.org/10.3920/WMJ2019.2449>
 44. Moghaddam AF, Rychlik M, Hosseini H, Janat B, Yazdanpanah H, et al. 2019. Risk associated with the intake of aflatoxin M1 from milk in Iran. *World Mycotoxin J* 12(2): 191-200. <https://doi.org/10.3920/WMJ2018.2368>
 45. Hussein HS, Brasel JM. 2001. Toxicity, metabolism, and impact of mycotoxins on humans and animals. *Toxicology* 167(2): 101-134. [https://doi.org/10.1016/S0300-483X\(01\)00471-1](https://doi.org/10.1016/S0300-483X(01)00471-1)
 46. Benites AJ, Fernandes M, Boleto AR, Azevedo S, Silva S, et al. 2017. Occurrence of ochratoxin A in roasted coffee samples commercialized in Portugal. *Food Control* 73: 1223-1228. <https://doi.org/10.1016/j.foodcont.2016.10.037>
 47. Pierron A, Alassane-Kpembé I, Oswald IP. 2016. Impact of mycotoxin on immune response and consequences for pig health. *Anim Nutr* 2(2): 63-68. <https://doi.org/10.1016/j.aninu.2016.03.001>
 48. Norbäck D, Hashim JH, Cai GH, Hashim Z, Ali F, et al. 2016. Rhinitis, ocular, throat and dermal symptoms, headache and tiredness among students in schools from Johor Bahru, Malaysia: associations with fungal DNA and mycotoxins in classroom dust. *PLoS One* 11(2): e0147996. <https://doi.org/10.1371/journal.pone.0147996>
 49. Schütze N, Lehmann I, Bönisch U, Simon JC, Polte T. 2010. Exposure to mycotoxins increases the allergic immune response in a murine asthma model. *Am J Respir Crit Care Med* 181(11): 1188-1199. <https://doi.org/10.1164/rccm.200909-1350OC>
 50. Selamat J, Iqbal SZ. 2016. Food Safety: Basic Concepts, Recent Issues, and Future Challenges. Springer.
 51. Puangkham S, Poapolathep A, Jermnak U, Imsilp K, Tanhan P, et al. 2017. Monitoring and health risk of mycotoxins in imported wines and beers consumed in Thailand. *World Mycotoxin J* 10(4): 401-409. <https://doi.org/10.3920/WMJ2017.2216>
 52. Ehrlich KC, Moore GG, Mellon JE, Bhatnagar D. 2015. Challenges facing the biological control strategy for eliminating aflatoxin contamination. *World Mycotoxin J* 8(2): 225-233. <https://doi.org/10.3920/WMJ2014.1696>
 53. Abrar M, Anjum FM, Butt MS, Pasha I, Randhawa MA, et al. 2013. Aflatoxins: biosynthesis, occurrence, toxicity, and remedies. *Crit Rev Food Sci Nutr* 53(8): 862-874. <https://doi.org/10.1080/10408398.2011.563154>
 54. Ramírez-Camejo LA, Zuluaga-Montero A, Lázaro-Escudero M, Hernández-Kendall V, Bayman P. 2012. Phylogeography of the cosmopolitan fungus *Aspergillus flavus*: is everything everywhere? *Fungal Biol* 116(3): 452-463. <https://doi.org/10.1016/j.funbio.2012.01.006>
 55. Tahir NI, Hussain S, Javed M, Rehman H, Shahzady TG, et al. 2018. Nature of aflatoxins: their extraction, analysis, and control. *J Food Saf* 38(6): e12561. <https://doi.org/10.1111/jfs.12561>
 56. Ostry V, Malir F, Toman J, Grosse Y. 2017. Mycotoxins as human carcinogens—the IARC monographs classification. *Mycotoxin Res* 33: 65-73. <https://doi.org/10.1007/s12550-016-0265-7>
 57. Ismail A, Gonçalves BL, de Neeff DV, Ponzilacqua B, Coppa CF, et al. 2018. Aflatoxin in foodstuffs: occurrence and recent advances in decontamination. *Food Res Int* 113: 74-85. <https://doi.org/10.1016/j.foodres.2018.06.067>
 58. Safety Evaluation of Certain Contaminants in Food: Prepared by the Eighty-third Meeting of the Joint FAO/WHO Expert Committee on Food Additives (JECFA). [<https://www.who.int/publications/item/9789240060760>] [Accessed November 05, 2023]
 59. Piotrowska M, Slizewska K, Biernasiak J. 2013. Mycotoxins in Cereal and Soybean-based Food and Feed. In El-Shemy H (ed) Brazil: Soybean-Pest Resistance. IntechOpen, pp 185-230.
 60. Voss KA, Gelineau-van Waes JB, Riley RT. 2006. Fumonisin: current research trends in developmental toxicology. *Mycotoxin Res* 22: 61-69. <https://doi.org/10.1007/BF02954559>
 61. Akbar A, Majeed FA, Sadiq MB, Khan SA, Rabaan AA. 2022. Mycotoxins occurrence in food commodities, their associated hazards and control strategies. *J Hellenic Vet Med Soc* 73(1): 3853-3866. <https://doi.org/10.12681/jhvms.26023>

62. Ben Hassouna K, Ben Salah-Abbes J, Chaieb K, Abbès S. 2022. Mycotoxins occurrence in milk and cereals in North African countries—a review. *Crit Rev Toxicol* 52(8): 619-635. <https://doi.org/10.1080/10408444.2022.2157703>
63. Alberts JF, Lilly M, Rheeder JP, Burger HM, Shephard GS, et al. 2017. Technological and community-based methods to reduce mycotoxin exposure. *Food Control* 73: 101-109. <https://doi.org/10.1016/j.foodcont.2016.05.029>
64. Claeys L, Romano C, De Ruyck K, Wilson H, Fervers B, et al. 2020. Mycotoxin exposure and human cancer risk: a systematic review of epidemiological studies. *Compr Rev Food Sci Food Saf* 19(4): 1449-1464. <https://doi.org/10.1111/1541-4337.12567>
65. Ostry V, Malir F, Ruprich J. 2013. Producers and important dietary sources of ochratoxin A and citrinin. *Toxins* 5(9): 1574-1586. <https://doi.org/10.3390/toxins5091574>
66. Palencia ER, Hinton DM, Bacon CW. 2010. The black *Aspergillus* species of maize and peanuts and their potential for mycotoxin production. *Toxins* 2(4): 399-416. <https://doi.org/10.3390/toxins2040399>
67. Bhat R, Rai RV, Karim AA. 2010. Mycotoxins in food and feed: present status and future concerns. *Compr Rev Food Sci Food Saf* 9(1): 57-81. <https://doi.org/10.1111/j.1541-4337.2009.00094.x>
68. Wan J, Chen B, Rao J. 2020. Occurrence and preventive strategies to control mycotoxins in cereal-based food. *Compr Rev Food Sci Food Saf* 19(3): 928-953. <https://doi.org/10.1111/1541-4337.12546>
69. Berthiller F, Crews C, Dall'Asta C, Saeger SD, Haesaert G, et al. 2013. Masked mycotoxins: a review. *Mol Nutr Food Res* 57(1): 165-186. <https://doi.org/10.1002/mnfr.201100764>
70. Bushnell WR, Perkins-Veazie P, Russo VM, Collins J, Seeland TM. 2010. Effects of deoxynivalenol on content of chloroplast pigments in barley leaf tissues. *Phytopathology* 100(1): 33-41. <https://doi.org/10.1094/PHYTO-100-1-0033>
71. Foroud NA, Shank RA, Kiss D, Eudes F, Hazendonk P. 2016. Solvent and water mediated structural variations in deoxynivalenol and their potential implications on the disruption of ribosomal function. *Front Microbiol* 7: 1239. <https://doi.org/10.3389/fmicb.2016.01239>
72. Holanda DM, Kim SW. 2021. Investigation of the efficacy of mycotoxin-detoxifying additive on health and growth of newly-weaned pigs under deoxynivalenol challenges. *Anim Biosci* 34(3): 405. <https://doi.org/10.5713/ajas.20.0567>
73. Subak H, Selvolini G, Macchiagodena M, Ozkan-Ariksoysal D, Pagliai M, et al. 2021. Mycotoxins aptasensing: from molecular docking to electrochemical detection of deoxynivalenol. *Bioelectrochemistry* 138: 107691. <https://doi.org/10.1016/j.bioelechem.2020.107691>
74. Lai Y, Sun M, He Y, Lei J, Han Y, et al. 2022. Mycotoxins binder supplementation alleviates aflatoxin B1 toxic effects on the immune response and intestinal barrier function in broilers. *Poult Sci* 101(3): 101683. <https://doi.org/10.1016/j.psj.2021.101683>
75. El Khoury D, Fayjaloun S, Nassar M, Sahakian J, Aad PY. 2019. Updates on the effect of mycotoxins on male reproductive efficiency in mammals. *Toxins* 11(9): 515. <https://doi.org/10.3390/toxins11090515>
76. Sobrova P, Adam V, Vasatkova A, Beklova M, Zeman L, et al. 2010. Deoxynivalenol and its toxicity. *Interdiscip Toxicol* 3(3): 94-99. <https://doi.org/10.2478/v10102-010-0019-x>
77. Janik E, Niemcewicz M, Podogrocki M, Ceremuga M, Stela M, et al. 2021. T-2 toxin—the most toxic trichothecene mycotoxin: metabolism, toxicity, and decontamination strategies. *Molecules* 26(22): 6868. <https://doi.org/10.3390/molecules26226868>
78. Nestic K, Ivanovic S, Nestic V. 2013. Fusarial Toxins: Secondary Metabolites of *Fusarium* Fungi. In Whitacre D (ed) *Reviews of Environmental Contamination and Toxicology* Volume 228. Springer, Cham, pp 101-120.
79. Mally A, Solfrizzo M, Degen GH. 2016. Biomonitoring of the mycotoxin zearalenone: current state-of-the art and application to human exposure assessment. *Arch Toxicol* 90: 1281-1292. <https://doi.org/10.1007/s00204-016-1704-0>
80. Nahle S, El Khoury A, Atoui A. 2021. Current status on the molecular biology of zearalenone: its biosynthesis and molecular detection of zearalenone producing *Fusarium* species. *Eur J Plant Pathol* 159(2): 247-258. <https://doi.org/10.1007/s10658-020-02173-9>
81. Huffman J, Gerber R, Du L. 2010. Recent advancements in the biosynthetic mechanisms for polyketide-derived mycotoxins. *Biopolymers* 93(9): 764-776. <https://doi.org/10.1002/bip.21483>
82. Alshannaq A, Yu JH. 2017. Occurrence, toxicity, and analysis of major mycotoxins in food. *Int J Environ Res Public Health* 14(6): 632. <https://doi.org/10.3390/ijerph14060632>
83. Pinotti L, Ottoboni M, Giromini C, Dell'Orto V, Cheli F. 2016. Mycotoxin contamination in the EU feed supply chain: a focus on cereal byproducts. *Toxins* 8(2): 45. <https://doi.org/10.3390/toxins8020045>
84. da Rocha MEB, Freire FDCO, Maia FEF, Guedes MIF, Rondina D. 2014. Mycotoxins and their effects on human and animal health. *Food Control* 36(1): 159-165. <https://doi.org/10.1016/j.foodcont.2013.08.021>
85. Wright SA. 2015. Patulin in food. *Curr Opin Food Sci* 5: 105-109. <https://doi.org/10.1016/j.cofs.2015.10.003>
86. Marin S, Ramos AJ, Cano-Sancho G, Sanchis V. 2013. Mycotoxins: occurrence, toxicology, and exposure assessment. *Food Chem Toxicol* 60: 218-237. <https://doi.org/10.1016/j.fct.2013.07.047>
87. Ráduly Z, Szabó L, Madar A, Pócsi I, Csernoch L. 2020. Toxicological and medical aspects of *Aspergillus*-derived mycotoxins entering the feed and food chain. *Front Microbiol* 10: 2908. <https://doi.org/10.3389/fmicb.2019.02908>
88. Zhong L, Carere J, Lu Z, Lu F, Zhou T. 2018. Patulin in apples and apple-based food products: the burdens and the mitigation strategies. *Toxins* 10(11): 475. <https://doi.org/10.3390/toxins10110475>
89. Plotka-Wasyłka J. 2018. A new tool for the evaluation of the analytical procedure: green analytical procedure index. *Talanta* 181: 204-209. <https://doi.org/10.1016/j.talanta.2018.01.013>
90. Han Z, Ren Y, Liu X, Luan L, Wu Y. 2010. A reliable isotope dilution method for simultaneous determination of fumonisins B1, B2 and B3 in traditional Chinese medicines by ultra-high-performance liquid chromatography-tandem mass spectrometry. *J Sep Sci* 33(17-18): 2723-2733. <https://doi.org/10.1002/jssc.201000423>
91. Han Z, Ren Y, Zhou H, Luan L, Cai Z, et al. 2011. A rapid method for simultaneous determination of zearalenone, α -zearalenol, β -zearalenol, zearalanone, α -zearalanol and β -zearalanol in traditional Chinese medicines by ultra-high-performance liquid chromatography-tandem mass spectrometry. *J Chromatogr B* 879(5-6): 411-420. <https://doi.org/10.1016/j.jchromb.2010.12.028>
92. Cho HD, Suh JH, Feng S, Eom T, Kim J, et al. 2019. Comprehensive analysis of multi-class mycotoxins in twenty different species of functional and medicinal herbs using liquid chromatography-tandem mass spectrometry. *Food Control* 96: 517-526. <https://doi.org/10.1016/j.foodcont.2018.10.007>
93. Gałuszka A, Migaszewski ZM, Konieczka P, Namieśnik J. 2012. Analytical eco-scale for assessing the greenness of analytical procedures. *Trends Anal Chem* 37: 61-72. <https://doi.org/10.1016/j.trac.2012.03.013>
94. Wang Y, Zhang C, Zhang L, Zhao M, OuYang Z, et al. 2020. A comprehensive strategy for screening and exploring multi-class mycotoxins contamination status in Astragali Radix. *Microchem J* 158: 105294. <https://doi.org/10.1016/j.microc.2020.105294>
95. Pena-Pereira F, Wojnowski W, Tobiszewski M. 2020. AGREE—Analytical GREENness metric approach and software. *Anal Chem* 92(14): 10076-10082. <https://doi.org/10.1021/acs.analchem.0c01887>
96. Shanakhat H, Sorrentino A, Raiola A, Romano A, Masi P, et al. 2018. Current methods for mycotoxins analysis and innovative strategies for their reduction in cereals: an overview. *J Sci Food Agric* 98(11): 4003-4013. <https://doi.org/10.1002/jsfa.8933>

97. Zhang Z, Hu X, Zhang Q, Li P. 2016. Determination for multiple mycotoxins in agricultural products using HPLC-MS/MS via a multiple antibody immunoaffinity column. *J Chromatogr B* 1021: 145-152. <https://doi.org/10.1016/j.jchromb.2016.02.035>
98. Gbashi S, Njobeh PB, De Saeger S, De Boevre M, Madala NE. 2020. Development, chemometric-assisted optimization and in-house validation of a modified pressurized hot water extraction methodology for multi-mycotoxins in maize. *Food Chem* 307: 125526. <https://doi.org/10.1016/j.foodchem.2019.125526>